



MANAGING PAIN FOR YOUR PRACTICE



Injectable Local Anaesthetics

SEPTANEST 1:100,000

4% articaine with 1:100,000 epinephrine injection solution

Indications

Local or loco-regional dental anaesthesia in patients of at least 4 years old

Features and Benefits

- Septanest 40 mg/ml plus adrenaline 0.01 mg/ml (1/100 000) is particularly appropriate for procedures of longer duration and when there is a risk of significant bleeding into the operative field
- Fast onset to avoid delay before effective analgesia is achieved
- No-methylparaben formulation to avoid allergic reactions on sensitive patients
- 100% latex-free
- Terminal sterilization of product/cartridges, the Gold Standard method for Health Authorities around the world
- Mylar wrapped cartridges to minimize risk of injury in the event of a cartridge breakage with blister packaging to avoid contamination

Presentation

- 2 x Box of 50 2.2ml cartridges



SEPTANEST 1:200,000

4% articaine with 1:200,000 epinephrine injection solution

Indications

Local or loco-regional dental anaesthesia in patients of at least 4 years old

Features and Benefits

- Septanest containing 1/200 000 epinephrine should be preferred as it reduces the risk of excessive uptake of Septanest into the systemic circulation
- No-methylparaben formulation to avoid allergic reactions on sensitive patients
- 100% latex-free
- Terminal sterilization of product/cartridges, the Gold Standard method for Health Authorities around the world
- Mylar wrapped cartridges to minimize risk of injury in the event of a cartridge breakage with blister packaging to avoid contamination

Presentation

- 2 x Box of 50 2.2ml cartridges

